

Section 7 - Literature Cited

- Andrews, J.D. 1988. Epizootiology of the disease caused by oyster pathogen, *Perkinsus marinus*, and its effect on the oyster industry. American Fisheries Society Special Publication 18:47-63.
- Batiuk, R.A., R.J. Orth, K.A. Moore, W.C. Dennison, J.C. Stevenson, L.W. Staver, V. Carter, N.B. Rybicki, R.E. Hickman, S. Kollar, S. Bieber, and P. Heasley 1992. Chesapeake Bay submerged aquatic vegetation habitat requirements and restoration targets: A technical synthesis. Chesapeake Bay Program CBP/TRS 83/92.
- Beck, M. W., K.L. Heck, K. W. Able, D. L. Childers, D.B. Eggleston, B.M. Gillanders, B. Halpern, C. G. Hays, K. Hoshino, T. J. Minello, R. J. Orth, P.F. Sheridan and M. P. Weinstein. 2001. The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *BioScience* 51: 633-641.
- Bortone, S. A. and R. K. Turpin. 2000. Tapegrass life history metrics associated with environmental variables in a controlled estuary. p. 65-79. In S.A. Bortone (ed.), Seagrass monitoring, ecology, physiology and management. C. R. C. Press, Boca Raton, Florida.
- Brock, D.A. 2001. Nitrogen budget for low and high freshwater inflows, Nueces Estuary, Texas. *Estuaries* 24: 509-521.
- Browder, J.A. and D. Moore 1981. A new approach to determining the quantitative relationship between fishery production and the flow of fresh water to estuaries. p. 403-430 In R.D. Cross and D.L. Williams (eds.), Proceedings of the national symposium on freshwater inflow to estuaries Volume 1. U.S. Fish and Wildlife Service, U.S. Dept. of Interior. FWS/OBS-81/04.
- Burreson, E. M. and L.M. Ragone-Calvo 1996. Epizootiology of *Perkinsus marinus* disease of oysters in Chesapeake Bay with emphasis on data since 1985. *Journal of Shellfish Research* 15: 17-34.
- Carter, V., J. W. Barko, G. L. Godshalk, and N. B. Rybicki. 1988. Effects of submersed macrophytes on water quality in the tidal Potomac River, Maryland. *Journal of Freshwater Ecology* 4:493-501.
- Chamberlain, R.H. and P.H. Doering. 1998. Preliminary estimate of optimum freshwater inflow to the Caloosahatchee Estuary: A resource-based approach. p. 121- 130. In S.F. Treat (ed.), Proceedings of the Charlotte Harbor Public Conference and Technical Symposium; 1997 March 15-16; Punta Gorda, FL. Charlotte Harbor National Estuary Program Technical Report No. 98-02.
- Chu, F.L.E. and A.K. Volety. 1997. Disease processes of the parasite *Perkinsus marinus* in the eastern oyster *Crassostrea virginica*: minimum dose for infection initiation, and interaction of temperature, salinity and cell dose. *Diseases of Aquatic Organisms* 28: 61-68.
- Dennison, W.C., R.J. Orth, K.A. Moore, J.C. Stevenson, V. Carter, S. Kollar, P. W. Bergstrom and R.A. Batiuk. 1993. Assessing water quality with submersed aquatic vegetation. *Bioscience* 43:86-94.

- Doering, P.H. and R.H. Chamberlain 1999. Water quality and the source of freshwater discharge to the Caloosahatchee Estuary, FL. Water Resources Bulletin 35: 793-806.
- Doering, P. H., R. H. Chamberlain and D. E. Haunert 2002. Using submerged aquatic vegetation to establish minimum and maximum freshwater inflows to the Caloosahatchee Estuary, FL. *Estuaries*. In press.
- Doering, P.H. and R.H. Chamberlain 2000. Experimental studies on the salinity tolerance of turtle grass *Thalassia testudinum*. Ch 6. pp 81-97, In S.A. Bortone (ed), *Seagrass: Monitoring ecology, physiology, and management*. CRC Press (Boca Raton, FL) 318 pp.
- Doering, P.H., R.H. Chamberlain, K.M. Donohue, and A.D. Steinman 1999. Effect of salinity on the growth of *Vallisneria americana* Michx. from the Caloosahatchee Estuary, Florida. *Florida Scientist* 62(2): 89-105.
- Doering, P.H. and R.H. Chamberlain 1998. Water quality in the Caloosahatchee Estuary, San Carlos Bay and Pine Island Sound. Proceedings of the Charlotte Harbor Public Conference and Technical Symposium; 1997 March 15-16; Punta Gorda, FL. Charlotte Harbor National Estuary Program Technical Report No. 98-02. 274 p.
- Edwards, R. E., W. Lung, P.A. Montagna, and H. L. Windom. 2000. Final review report. Caloosahatchee Minimum Flow Peer Review Panel, September 27-29, 2000. Report to the South Florida Water Management District, West Palm Beach, FL.
- Estevez, E.D. 2000a. Matching salinity metrics to estuarine seagrasses for freshwater inflow management. p. 295-307. In S. A. Bortone (ed.), *Seagrasses: Monitoring, ecology, physiology and management*. CRC Press, Boca Raton FL
- Flaig, E. G. and J. Capece 1998. Water use and runoff in the Caloosahatchee watershed. p. 73- 80 In S. F. Treat (ed.), *Proceedings of the Charlotte Harbor Public Conference and Technical Symposium; 1997 March 15-16; Punta Gorda, FL*. Charlotte Harbor National Estuary Program Technical Report No. 98-02
- Fonseca, M.S. and J.S. Fisher 1986. A comparison of canopy friction and sediment movement between four species of seagrass and with reference to their ecology and restoration. Marine Ecology Progress Series 29:15-22.
- Gunter, G. 1961. Some relations of estuarine organisms to salinity. Limnology and Oceanography 6: 182-190.
- Hopkinson, C.S., Jr. and J.J. Vallino. 1995. The relationships among man's activities in watersheds and estuaries: A model of runoff effects on patterns of estuarine community metabolism. Estuaries 18: 598-621.
- Jassby, A.D. , W.J. Kimmerer, S.G. Monismith, C. Armor, J.E. Cloern, T. M. Powell, J. R. Schubel and T.J. Vendlinski. 1995. Isohaline position as a habitat indicator for estuarine populations. Ecological Applications 5: 272-289.
- Johansson, J.O.R. and H. S. Greening. 2000. Seagrass restoration in Tampa Bay: A resource-based approach to estuarine management. p. 279-294. In S.A.Bortone (ed.), *Seagrasses: Monitoring, ecology, physiology and management*. CRC Press, Boca Raton FL.

- Kemp, W.M., W. R. Boynton, R. R. Twilley, J. C. Stevenson, and L. G. Ward. 1984. Influences of submersed vascular plants on ecological processes in upper Chesapeake Bay. p. 367-394 In V.S. Kennedy (ed.), *The estuary as a filter*. Academic Press, New York, N.Y.
- Killgore, K. J., R. P. Morgan II, and N. B. Rybicki. 1989. Distribution and abundance of fishes associated with submersed aquatic plants in the Potomac River. *North American Journal of Fisheries Management* 9:101-111.
- Kimmerer, W. J. and J. R. Shubel. 1994. Managing freshwater flows into San Francisco Bay using a salinity standard: results of a workshop. p. 411-416. In K. R. Dyer and R. J. Orth (eds.), *Changes in fluxes in estuaries: implications from science to management*. Olsen and Olsen, Fredensbourg.
- Kraemer, G.P., R.H. Chamberlain, P.H. Doering, A.D. Steinman and M.D. Hanisak. 1999. Physiological responses of *Vallisneria americana* transplants along a salinity gradient in the Caloosahatchee Estuary (SW Florida). *Estuaries* 22:138-148.
- Livingston, R.J., X. Niu, F.G. Lewis III, and G.C. Woodsum. 1997. Freshwater input to a Gulf Estuary: Long term control of trophic organization. *Ecological Applications* 7:277-299
- Longley, W. L. (ed.) 1994. Freshwater inflows to Texas bays and estuaries: Ecological relationships and methods for determination of needs. Texas Water development Board and Texas Parks and Wildlife D, Austin, TX.
- Lubbers, L., W. R. Boynton, and W. M. Kemp. 1990. Variations in structure of estuarine fish communities in relation to abundance of submersed vascular plants. *Marine Ecology Progress Series* 65: 1-14.
- Matsumodo, J., G. Powell and D. Brock. 1994. Freshwater – inflow need of estuary computed by Texas Estuarine MP model. *Journal of Water Resources Planning and Management* 120:693-714.
- Nixon, S.W., J.W. Ammerman, L.P. Atkinson, V.M. Berounsky, G. Billen, W.C. Boicourt, W.R. Boynton, T.M. Church, D.M. DiToro, R. Elmgren, J.H. Garber, A.E. Giblin, R.A. Jahnke, N.J.P. Owens, M.E.Q. Pilson and S.P. Seitzinger 1996. The fate of nitrogen and phosphorus at the land-sea margin of the North Atlantic Ocean. *Biogeochemistry* 35: 141-180.
- Peterson, M., 2002 (in review): Tidal Caloosahatchee Basin Model, DHI Water & Environment, Technical report to South Florida Water Management District.
- Postel, S.L., G.C. Daily, and P.R. Ehrlich 1996. Human appropriation of renewable fresh water. *Science* 271: 785-788.
- Powell, G. L. and J. Matsumodo. 1994. Texas mathematical programming model: a tool for freshwater inflow management. p. 401-404 In K. R. Dyer and R. J. Orth (eds.), *Changes in fluxes in estuaries: implications from science to management*. Olsen and Olsen, Fredensbourg.
- Ray, S. M. 1954. Biological studies of *Derocystidium marinum*. The Rice Institute Pamphlet. Special Issue. November 1954, pp 65 – 76.

- Schlacher, T.A. and T.H. Wooldridge 1996. Ecological responses to reductions in freshwater supply and quality in South Africa's estuaries: lessons for management and conservation. Journal of Coastal Conservation 2: 115-130.
- Sellers, M.A. and J.G. Stanley. 1984. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (North Atlantic) -- American Oyster. U.S. Fish and Wildlife Service FWS/OBS-82/11.23. U.S. Army Corps of Engineers, TR EL--82-4. 15 pp.
- Sheng, Y. P., 2001: Impact of Caloosahatchee Flow on Circulation & Salinity in Charlotte Harbor, Technical Report for South Florida Water Management District, Civil & Coastal Engineering Department, University of Florida, Gainesville, Florida
- Sklar, F.H. and J. A. Browder 1998. Coastal environmental impacts brought about by alterations to freshwater flow in the Gulf of Mexico. Environmental Management 22: 547-562.
- Soniat, T. M. 1996. Epizootiology o Perkinsu marinus disease of eastern oysters in the Gulf of Mexico. Journal of Shellfish Research 15: 35-43.
- South Florida Water Management District. 2000b. Lower West Coast Water Supply Plan, Water Supply Department, Water Resources Management, SFWMD, West Palm Beach, FL.
- Stevenson, J. C., L.W. Staver and K.W. Staver. 1993. Water quality associated with survival of submersed aquatic vegetation along an estuarine gradient. Estuaries 16: 346-361.
- Tomasko, D.A. , C.J. Dawes and M.O. Hall. 1996. The effects of anthropogenic enrichment on Turtle Grass (Thalassia testudinum) in Sarasota Bay, Florida. Estuaries 19 (2B): 448-456.
- Thayer, G. W., W.J. Kenworthy, and M.S. Fonseca 1984. The ecology of eelgrass meadows of the Atlantic Coast: A community profile. U.S. Fish and Wildlife Service Report No. FWS/OBS-84/02 147 pp.
- United States Army Corps of Engineers and South Florida Water Management District, 1999. The Central and Southern Florida Flood Control Project Comprehensive Review Study. USACE, Jacksonville District, Jacksonville, FL. and SFWMD, West Palm Beach, FL.
- Virnstien, R.W. and L.J. Morris 2000. Setting seagrass targets for the Indian River Lagoon, Florida. p. 211-218. in S. A. Bortone (eds), Seagrasses: Monitoring, ecology, physiology and management. CRC Press, Boca Raton FL. 318 p.
- Whitfield, A.K. and T.H. Wooldridge 1994. Changes in freshwater supplies to southern African estuaries: some theoretical and practical considerations. p. 41-50. In K. R. Dyer and R. J. Orth (eds.), Changes in fluxes in estuaries: Implications from science and management. Olsen and Olsen, Fredensborg
- Zieman, J.C., and R.T Zieman. 1989. The ecology of the seagrass meadows of the west coast of Florida: a community profile. U. S. Fish and Wildlife Service Biological Report 85(7.25).